UGC Approved Indexed Peer Reviewed Journal

ISSN: 2320-6586 (Online)

IILM UNIVERSITY

Technologies for Sustainable Development Goals 2025 (TECH4SDG-2025)

School of Computer Science & Engineering

IILM University Greater Noida

Presents

1st National Conference

on

Technologies for Sustainable Development Goals 2025 (TECH4SDG-2025)

3rd March 2025

Technically Co-Sponsored by IEEE UP SECTION (INDIA)

Venue:

IILM University, Greater Noida

Plot No.18,16, Knowledge Park II, Greater Noida, Uttar Pradesh 201306

Procedings Published at

International Journal of Research in Modern Engineering and Emerging Technology

Vol. 13, Sp. Issue 1, April 2025

Doi: https://doi.org/10.63345/ijrmeet.org.v13.si1.1301

Editor-in-Chief

Dr. Mylavarapu Seetharama Prasad

www.ijrmeet.org

INDEX

S. No.	Particulars	Page No.
1	A Comparative Study On Load Balancing Algorithms And Its Types **Methily Johri & Aditri Ashish**	1–10
2	Optimizing Supply Chain Management with Ethereum 2.0: Innovations in Blockchain Technology **Ayush Gupta & Dr. Ambuj Kumar Agarwal**	11–17
3	Smart Surveillance: "A Real-Time Crowd Analysis System for Density and Anomaly Detection" **Komal Sharma, Dr. Pankaj Kumar Varshney, Aditi Aggarwal & Gaurav Kumar**	17–22
4	AI-Driven Network Optimization: Matching Policyholders to Cost-Effective Providers Using Business Intelligence **Ruchi Mangharamani & Prof. (Dr) Sangeet Vashishtha**	23–29
5	The Future of UX: Designing for Emerging Technologies (AI, AR, VR, Voice UI) **Ojas Gupta & Prof.(Dr.) Arpit Jain**	30–37
6	Enhancing Liquidity Stress Testing: Best Practices for Modern Banking Risks **Padmini Bulani & Dr Abhishek Jain**	38–45
7	AI-Powered Investment Strategies: Optimizing Risk and Returns in Modern Banking **Aditya Mehra & Dr. Shakeb Khan**	46–53
8	Scaling Customer Journey Optimization in a Multi-Tenant Distributed System **Sumit Kumar Agrawal & Prof. (Dr) MSR Prasad**	54–61
9	Object Detection Using Convolutional Neural Networks: Advances, Techniques, and Applications	62–69

S. No.	Particulars	Page No.
	Bharath Thandalam Rajasekaran, Saravanan Thirumazhisai Prabhagaran & Prof. (Dr) Punit Goel	
10	AI-Augmented Intrusion Detection Systems for High-Speed Optical Networks **Sattvik Sharma & Prof.(Dr) Avneesh Kumar**	70–77
11	Generative AI in Data Security: Opportunities and Challenges **Vybhavreddy Kammireddy Changalreddy, Piyush Bipinkumar Desai & Prof (Dr) Ajay Shriram Kushwaha**	78–86
12	Integrating Salesforce with Cloud-Based Data Warehousing for Unified Analytics **Nidhi Shashikumar & Dr S P Singh**	87–94
13	AI-Powered Cybersecurity: A Unified Approach to Protecting Enterprise, Cloud, and SaaS Applications **Varun Grover, Khushmeet Singh & Er. Siddharth**	95–103
14	Sales and Order Analysis Using Excel-Based Dashboard: A Case Study on Vrinda Store **Mohd Ammar Khan & Harun Faridi**	104–106
15	ChatMania: A Realtime Chat Application **Dr. Kirti Shukla & Anshika**	107–110
16	Digital Surveillance and Domestic Abuse: A Legal Vacuum in Indian Marital Relationships **Vishal Nehra & Dr. Rakesh Kumar**	110–116
17	The Role of Emotional Appeal in Online Advertisements Targeting Working Women in Delhi-NCR **Mona Mishra & Dr. Praveen Mittal**	116–120
18	AI-Driven Data Governance Strategies for Enhanced Cloud Security in Enterprises **Chandana Pandey & Dr Gauri Shanker Kushwaha**	121–123

S. No.	Particulars	Page No.
19	Evolution of Social Media Advertising and Consumer Behavior **Mr. Pramod Rana & Prof (Dr) Punit Goel**	124–129
20	Sustainability Accounting Practices in Indian Manufacturing Firms **Mrs. Sonal Jain & Dr. Ajit Singh**	129–134
21	Fundamental Rights vs. Reasonable Restrictions: A Study of Evolving Judicial Balances **Mrs. Anshika Agarwal & Dr. Vijay Kumar Singh**	134–139
22	सतत शिक्षा के लिए हिंदी साहित्य की भूमिका: सतत विकास लक्ष्य-4 (गुणवत्तापूर्ण शिक्षा) का सांस्कृतिक विश्लेषण **श्रीमती रेणु गुप्ता, डॉ. बृजलता शर्मा**	140–144
23	Mind–Gut Synergy in Chronic Lifestyle Conditions: Evaluating Psychotherapy-Driven Behavioral Change and Microbiota-Mediated Metabolic Repair **Krithika. V, Dr. Sarbesh Kumar Singh**	145–151
24	प्राचीन शिक्षा प्रणाली से आधुनिक AI आधारित शिक्षा की ओर संक्रमण **गौरव जोशी, डॉ सारिका गोयल**	152–159
25	पर्वतीय क्षेत्रों में शिक्षा के डिजिटलीकरण की चुनौतियाँ एवं संभावनाएँ **राकेश उनियाल, डॉ. मीनाक्षी हर्राव वर्मा**	160–168
26	Women Leadership and Democratic Participation in India: Challenges and Opportunities in Parliamentary Politics **Renu Yadav, Dr. P.P.S. Rana**	169–176
27	Evaluating GST Effects on Competitiveness and Industrial Development **Aman Goel, Dr. Pooja Yadav**	177–183
28	एस०आर० हरनोट के कथा साहित्य में लोक संस्कृति, पर्यावरण चेतना और सामाजिक परिवर्तन का अध्ययन **सीमा देवी, डॉ० अरविन्द्र कौर चुंबर**	184–189

S. No.	Particulars	Page No.
29	Balancing National Security and Human Rights in Modern India **Shalu Jain, Dr. Vijay Kumar**	190–197
30	Economic Performance Drivers and Fiscal Resilience in Indian Airports **Lalit Kumar Sharma, Dr. Pooja Yadav**	198–203
31	Educational Governance and AI Learning Outcomes in Government Elementary Schools **Dr. Shubhra Chaturvedi**	203–210
32	Artificial Intelligence and the Future of Teacher Roles in India Classroom **Neeru Verma**	211–220
33	Research Culture Among Undergraduate Students in Indian Universities **Dr. Dhanashri Rajshri Ravindra Jadhav**	221–229
34	डिजिटल समाज में बहुभाषिक संचार का बदलता स्वरूप: भारतीय भाषाओं के समक्ष अवसर और चुनौतियाँ **Rakesh Uniyal**	230–234
35	Graduate Employability and Skill Gaps in State Universities **Narender Panghal**	235–242
36	Political Funding Transparency in Indian Election **Dr. Anand Kumar Dwivedi**	243–252
37	AI-Powered Platforms for Data Analytics and Audience Development **Sachin Sangal**	253–258
38	Engineering Leadership Models in AI Startups: Balancing Innovation and Operational Efficiency **Dr. Kalpesh Rasiklal Rakholia**	259–269
39	Challenges in Integrating CWSN in Regular Classrooms **Anil Kumar**	270–280
40	An Artificial Intelligence-Based Legal Compliance Framework for Real-Time Environmental Governance to Advance SDGs 9, 12, 13, and 16 **Dr. Jyoti Janardhan Reddy**	281–288

S. No.	Particulars	Page No.
41	Impact of Source Segregation and Waste Characterization on the Efficiency of Material Recovery Facilities **Varinder Vikram Singh, Dr. Pankaj Kumar Bharti**	289–296
42	Significance of Emotional Intelligence of schedule caste students in relation to Self-concept at secondary level **Mrs. Varsha Asher Wasker, Dr. Kuldeep Singh Tomar**	297–305
43	औपनिवेशिक भारत में स्थानीय बाजारों के सामाजिक परिवर्तन का ऐतिहासिक अध्ययन **Pankaj Tamta**	306–310
44	Teachers' Readiness to Integrate Generative AI into Lesson Planning and Classroom Instruction **Prema Bharti**	311–316
45	Battery Usage Optimization Techniques for AI-Heavy Mobile Apps **Dr. Akankasha Kathuria**	317–321
46	Gender, Technology, and Changing Family Structures in Urban India: A Socio-cultural Analysis **Prof. Sawant Jeetendra Madhukar Usha**	322–326
47	Law and Alternative Dispute Resolution in Contemporary Civil Justice: A Legal Study **Dr. Pooja Khurana**	327–331
48	English Literature and the Representation of Women in Contemporary Fiction: A Corpus-Based Study of Narrative Agency, 2000–2024 **Dr. Abu Farah Hoque**	332–336
49	Agricultural Impact of Biochar and Reduced Fertilizer on Mustard Soil Health and Yield **Dr. Bibhu Santosh Behera**	337–346
50	Ethical and Privacy Frameworks for AI-Driven Customer Data Analytics in Medical Profession **Dr. Gaurav Singh Tomar**	347–351
51	Legal Protection of Witnesses in Criminal Justice Proceedings **Dr. Jyoti Janardhan Reddy**	352–357

S. No.	Particulars	Page No.
52	Platform Culture and Fragmented Selfhood in Contemporary English Literature **Dr. Akanksha Singh**	358–362
53	Digital Health Literacy and Health Information Inequality in Rural India: A Multidimensional Assessment **Ranjana Jadaun**	363–368
54	स्मार्ट विनिर्माण और उद्योग 5.0 में कृत्रिम बुद्धिमत्ता, मानव-केंद्रित तकनीक और सतत विकास का अंतःविषयक अध्ययन **Dr. Ghamendra Kumar Sahu**	369–375
55	Contribution of Yoga-Based Breathing Techniques to Daily Stress Management **Dr. Sweta Kumari**	376–387
56	Algorithmic Silence and Unspoken Meaning in Contemporary English Fiction **Dr. Jyoti Devi**	388–396
57	Cost-Benefit Analysis of Custom Image Analysis Tools in Oncology Research **Sumit Malhotra**	397–405
58	उच्च शिक्षा में सूक्ष्म अधिगम तकनीकों का शैक्षिक प्रभाव **Ram Singh**	406–412
59	Optimising Laboratory Workflows with Low-Code/No-Code Software Solutions **Dr. Amit Nampalliwar**	413–421
60	भारतीय बहुभाषिक समाज में भाषा, संस्कृति और राष्ट्रीय एकता का अध्ययन **Dr. Geeta Rawat**	422–428
61	Influence of FinTech Innovation on Financial Resilience and Operational Efficiency of Retailers and Street Vendors **Prof. Manmeet Kumar Siras & Shyam Pandey**	429–435
62	भारत में आपातकालीन शक्तियाँ और लोकतांत्रिक उत्तरदायित्व: संसदीय शासन के अंतर्गत राष्ट्रपति की संवैधानिक भूमिका का विश्लेषण **सीता राम पंडित, डॉ. पी. पी. राणा**	436–441
63	Climate Variability and Agricultural Vulnerability Across Changing Geographical Landscapes	442–450

S. No.	Particulars	Page No.
	Dr. Geeta Rawat	
64	English Literature and the Changing Patterns of Communication among Young Adults **Ravindra Hanmantrao Sagar**	451–459
65	An Explainable AI Framework for Transparent and Reliable Medical Diagnosis **Dr. Thejoram Naresh Reddy Boya**	460–466
66	डिजिटल समुदायों का युवा शिक्षा और सामूहिक पहचान पर प्रभाव **डॉ. मनोज कुमार सकलानी**	467–473
67	Adaptive Technologies for Enhancing Online Shopping Experiences **Dr. Abhishek Bhattacharjee**	474–479
68	Impact of Digital Payment Adoption through AI-ML on Micro-Business Performance **Ralphy Joseph C J**	480–484
69	Role of Regional Parties in Shaping Federalism in India **Dr. Shailendra Kumar**	485–489
70	Serverless Computing Optimization: Enhancing Performance with AI-Based Resource Allocation **S. Venkatasubramanian**	490–495
71	Student Satisfaction in Autonomous vs. Affiliated Colleges **Dr. Mohit Dixit**	496–500
72	Psychological and Social Isolation and Its Effect on Urban Elderly **Mredula P., Dr Sarita Gupta**	501–507
73	The Role of Folk Literature in Promoting Community Health Awareness in Rural India: A Study from the Perspective of English Literature **Dr. Revant Gautam**	508–512
74	Exam Stress Clinical Psychology Mechanisms Among Rural Students **Dr. Avinash D. Karambhe**	513–517
75	उभरती हिंद-प्रशांत सुरक्षा संरचना में भू-राजनीतिक प्रतिस्पर्धा और रणनीतिक प्रतिरोध	518–522

S. No.	Particulars	Page No.
	Dr. Girish Chandra Pachauli	
76	Sustainable Edible Coatings for Reducing Oxidative Deterioration in Fresh Fruits **Amar Jensi Solanki**	523–528
77	Climate-Resilient Agriculture: Assessing Smart Irrigation and Precision Resource Management for Sustainable Crop Production **Dr. Bibhu Santosh Behera**	529–534

Contribution of Yoga-Based Breathing Techniques to Daily Stress Management

Dr. Sweta Kumari
TGT Yoga Teacher
Kendriya Vidyalaya, Sec 24, Noida
Orcid id: https://orcid.org/0009-0002-7378-1613

Abstract— Daily psychological stress is increasingly recognized as a persistent challenge affecting emotional stability, cognitive performance, sleep quality, and general well-being among adults. Yoga-based breathing techniques offer a low-cost, non-pharmacological approach through which individuals may voluntarily regulate respiratory rhythm and potentially modify physiological and psychological responses to everyday stressors. Although recent studies have examined breathwork, pranayama, and comprehensive yoga programs, relatively limited attention has been given to brief breathing practices performed independently within routine daily environments. The present study addresses this gap by conceptualizing a digitally supported framework for examining the relationship between breathing-practice adherence, perceived stress variation, respiratory characteristics, and autonomic indicators. The proposed research distinguishes immediate stress regulation from cumulative stress reduction and evaluates whether consistency of practice contributes more strongly to stress outcomes than session duration alone. A multimodal experimental framework integrating self-reported stress measures, breathing-session records, heart-rate-derived variables, and contextual daily observations is proposed for subsequent empirical evaluation.

Keywords— Yoga-Based Breathing, Pranayama, Daily Stress, Breath Regulation, Digital Health, Autonomic Regulation

Introduction

Stress is not confined to major traumatic events or clinical psychiatric conditions. For a substantial proportion of the adult population, stress emerges repeatedly through occupational demands, academic responsibilities, interpersonal interactions, financial uncertainty, time pressure, digital overload, and disruptions to established routines. These recurring demands may be individually modest but can accumulate across a day, producing measurable changes in affect, attention, sleep, perceived well-being, and physiological arousal. Consequently, contemporary stress-management research increasingly requires interventions that can be incorporated into ordinary life rather than approaches that depend exclusively on specialized facilities, lengthy therapeutic sessions, or continuous professional supervision.

Breathing regulation represents one of the most accessible behavioral mechanisms through which individuals can intentionally influence their immediate psychophysiological state. Unlike several physiological processes associated with stress, respiration is both automatically regulated and voluntarily modifiable. Yoga traditions have long incorporated systematic manipulation of breathing through practices collectively described as *pranayama*. Depending upon the technique, practitioners may alter breathing frequency, inspiratory and expiratory duration, pauses between respiratory phases, nostril use, or attentional focus. In contemporary health research, these techniques increasingly intersect with the broader scientific category of controlled breathwork.

Recent evidence provides reasonable grounds for investigating breathing-based interventions as stress-management tools. A meta-analysis of randomized controlled breathwork trials reported a statistically significant small-to-medium reduction in self-reported stress among participants receiving breathwork interventions compared with controls. Research examining yoga more broadly has similarly

indicated beneficial effects on perceived stress, although effect estimates vary considerably between interventions and study populations. A 2024 systematic review and meta-analysis involving stressed adults identified evidence favoring yoga for short-term stress reduction while simultaneously emphasizing limitations in the overall quality of available evidence.

More focused investigation of yoga-based respiration is particularly relevant because breathing exercises require comparatively little space, equipment, or physical mobility. They can potentially be practiced during a work break, before an examination, after commuting, between meetings, or before sleep. A person does not necessarily require the environmental conditions associated with a complete yoga session. This feature gives controlled breathing potential relevance to mobile health applications, wearable-device ecosystems, workplace wellness systems, and personalized digital interventions.

Evidence concerning acute effects is also developing. A systematic review examining single sessions of yoga, meditation, and breathing-related practices reported that a majority of eligible studies demonstrated outcomes favoring intervention conditions for either physiological or psychological stress reactivity. At the same time, the authors identified methodological limitations and called for stronger studies with more representative populations. These findings suggest that even relatively brief interventions may influence stress responses, but they do not establish that such effects translate reliably into the unsupervised circumstances of everyday life.

This distinction is important. An intervention that reduces stress in a laboratory, yoga center, or supervised session does not automatically become an effective daily stress-management strategy. Real-world effectiveness depends upon whether individuals actually perform the breathing exercise, whether the technique remains effective when practiced without an instructor, how frequently it must be repeated, and whether benefits persist beyond the immediate post-practice period. Adherence itself may also fluctuate according to stress: highly stressed individuals could have a greater need for breathing practice while simultaneously being less likely to perform it consistently.

Another unresolved issue concerns the dose-response characteristics of yoga-based breathing. Research protocols differ markedly in breathing frequency, session duration, intervention length, instructional format, and specific technique. For example, a randomized investigation of slow yoga-based breathing conducted over 12 weeks found reductions in psychological measures related to anxiety, while differences between breathing ratios were less clear and physiological outcomes did not necessarily parallel psychological changes. Such findings challenge the assumption that simply prescribing a particular respiratory ratio will generate predictable improvement across individuals.



Fig. 1: Multiple Benefits of Yoga as seen from many research studies.

Source: https://www.researchgate.net/figure/Multiple-Benefits-of-Yoga-as-seen-from-many-research-studies_fig1_393441966

The problem therefore requires analysis beyond the conventional question of whether breathing "works." A more useful research question is whether variation in practice characteristics—including frequency, adherence, timing, respiratory pace, and baseline stress—can explain differential outcomes. From a digital-health perspective, this shift is significant because modern wearable sensors and mobile applications can generate repeated observations of breathing behavior, pulse characteristics, practice timing, and

subjective stress. Such longitudinal data provide an opportunity to model stress regulation as a dynamic process rather than as a single pre-test/post-test comparison.

Recent research has also continued to examine pranayama in specialized populations. A 2025 systematic review and meta-analysis investigating pranayama among individuals with diagnosed mental disorders identified short-term improvements in symptom severity compared with passive controls, but emphasized the small evidence base and substantial concerns regarding risk of bias. A 2026 review of yoga-based voluntary breathing among medical students similarly reported beneficial pooled effects for stress and anxiety, while substantial heterogeneity and the small number of rigorous randomized trials limited generalization. These findings reinforce the therapeutic promise of breathing practices while also demonstrating why stronger designs are needed.

1.1 Research Gap

The principal gap addressed in the present manuscript concerns the **lack of ecologically grounded models explaining how adherence to brief yoga-based breathing practices interacts with momentary stress and physiological regulation during ordinary daily life**.

Much of the existing literature can be grouped into three categories: comprehensive yoga interventions combining postures, breathing, relaxation, or meditation; controlled breathing studies performed under supervised experimental conditions; and clinical or student-centered interventions evaluated predominantly through pre-intervention and post-intervention questionnaires. These approaches provide important evidence of efficacy but reveal comparatively little about day-to-day behavioral implementation.

In particular, limited research simultaneously evaluates:

1. how consistently participants perform short breathing sessions outside supervised environments;
2. whether stress immediately before practice influences the magnitude of post-practice change;
3. whether practice frequency is more important than individual session duration;
4. whether subjective stress improvements correspond with wearable-derived physiological responses; and
5. whether individual response patterns can be identified computationally rather than assuming a homogeneous treatment effect.

The present study therefore conceptualizes yoga-based breathing as a repeated behavioral micro-intervention rather than solely as a conventional yoga program. This distinction provides a more practical foundation for digital stress-management systems.

1.2 Research Objective

The central objective of the study is to evaluate the contribution of brief yoga-based breathing practices to daily stress regulation by integrating behavioral adherence, subjective stress measurements, respiratory characteristics, and physiological indicators within a longitudinal digital-health framework.

More specifically, the study is designed to determine whether repeated short breathing sessions are associated with measurable reductions in momentary and cumulative stress; to assess the relationship between practice regularity and outcome magnitude; to examine whether baseline stress modifies individual responsiveness; and to develop an interpretable predictive model capable of identifying patterns associated with stronger or weaker stress reduction.

Unlike conventional intervention studies relying exclusively on average pre-to-post differences, the proposed analysis treats each breathing session as an observation nested within a participant. This allows day-level and person-level effects to be examined separately and provides the basis for individualized stress-management recommendations.

Literature Review

2.1 Stress Regulation and the Respiratory System

Psychological stress involves coordinated cognitive, emotional, endocrine, and autonomic responses. Acute stress frequently produces changes in breathing rate, respiratory depth, cardiovascular activity, muscle tension, and attentional state. Respiratory patterns may themselves reinforce subjective arousal. Consequently, deliberate modification of

breathing has been investigated as a potential bottom-up mechanism for altering stress-related physiological activity.

Slow and controlled breathing is frequently associated with changes in autonomic regulation, although physiological responses depend on respiratory frequency, depth, individual characteristics, measurement methods, and experimental conditions. It is therefore overly simplistic to interpret every form of slow breathing as producing an identical parasympathetic response. The mechanisms are likely to involve interacting respiratory, cardiovascular, attentional, and interoceptive processes.

Yoga-based breathing provides a useful framework for studying these processes because it includes structured respiratory manipulations that can be standardized while remaining feasible for repeated self-practice.

2.2 Yoga and Perceived Stress

The wider yoga literature offers substantial contextual evidence for stress-management effects. A recent meta-analysis examining yoga and perceived stress reported an overall moderate effect while also identifying considerable heterogeneity across studies. This heterogeneity is scientifically important because it indicates that outcomes cannot be fully explained by the broad label "yoga."

Yoga interventions differ in physical intensity, meditative content, breathing content, instructor contact, treatment duration, population characteristics, and comparison conditions. Consequently, an intervention containing postures, relaxation, meditation, and breathing cannot determine how much improvement is attributable specifically to respiratory regulation.

A 2024 synthesis focusing on stressed adults likewise concluded that yoga may be considered a non-pharmacological option for stress reduction, but characterized the evidence for some stress outcomes as low quality. This supports further investigation while cautioning against exaggerated causal claims.

2.3 Pranayama as a Distinct Behavioral Intervention

Pranayama differs conceptually from generic relaxation because participants deliberately control components of the respiratory cycle. Depending on the practice, intervention characteristics may include slower breathing, prolonged expiration, alternate-nostril breathing, paced inhalation-exhalation ratios, breath observation, or brief respiratory pauses.

Isolating pranayama is methodologically valuable. When breathing is examined separately from yoga postures and meditation, researchers can more directly study its independent contribution to psychological outcomes. Recent systematic work supports continued investigation but reveals a relatively small and heterogeneous evidence base. For example, the 2025 meta-analysis of pranayama in mental disorders identified potential short-term benefits but also highlighted high risk of bias in several included trials.

An important safety consideration is that yoga breathing techniques are not physiologically interchangeable. The same review reported that adverse events were observed more frequently with fast breathing than with slow breathing practices. Accordingly, the proposed study emphasizes gentle, slow, non-forceful breathing rather than highly stimulating respiratory techniques.

2.4 Acute Versus Cumulative Effects

One of the most important distinctions in breathing research concerns acute and cumulative outcomes. An acute effect refers to the change occurring immediately after a breathing session—for example, a decline in self-rated stress within several minutes. A cumulative effect refers to longer-term change following repeated practice across days or weeks.

Existing studies sometimes blur these processes. Yet an intervention could produce a reliable immediate calming effect without substantially changing baseline stress over several weeks. Conversely, repeated breathing practice could gradually modify coping responses even when individual sessions produce modest immediate effects.

Evidence from single-session studies suggests that yoga components, including breathing-related practices, may reduce acute stress reactivity. Longitudinal slow-breathing research also suggests that repeated practice can influence psychological outcomes over extended periods. However, the relationship between the immediate and cumulative effects remains insufficiently characterized.

The proposed study therefore treats these outcomes separately. Momentary stress reduction will be assessed at the breathing-session level, whereas cumulative stress change will be analyzed across the entire intervention period.

2.5 Adherence as an Understudied Determinant

Behavioral interventions can only produce practical benefits when participants actually use them. Nevertheless, adherence is frequently reported merely as attendance or total completion rate rather than investigated as an explanatory variable.

For daily breathing practice, adherence is potentially multidimensional. It may involve the number of completed sessions, days practiced, regularity between sessions, duration of each session, and fidelity to the target breathing rate. Two individuals could therefore receive the same intervention but experience substantially different exposure.

This issue is especially relevant to digital health. Mobile applications can timestamp breathing sessions and record duration, while wearable sensors may provide complementary physiological data. Instead of treating adherence as a secondary implementation statistic, computational models can examine whether adherence predicts stress reduction and whether the relationship is linear, threshold-based, or person-specific.



Fig. 2: Breathing Meditation Yoga

Source: https://www.instagram.com/popular/breathing-meditation-yoga/

2.6 Digital Measurement and Personalized Stress Management

The expansion of wearable sensing and mobile self-monitoring has altered how behavioral interventions can be studied. Traditional studies often obtain stress questionnaires at baseline and after several weeks. Although useful for estimating overall intervention effects, this approach cannot capture substantial within-person variation occurring during the intervening days.

Repeated digital measurements provide a different data structure. A participant may contribute observations before and after multiple breathing sessions, allowing researchers to determine whether the intervention is more effective during high-stress periods, at particular times of day, or after repeated practice. Heart rate and heart-rate-variability-derived indicators can further provide physiological context, although such measures should not be interpreted as direct or exclusive measures of psychological stress.

Machine learning may be particularly useful when applied carefully to these longitudinal data. Rather than replacing conventional statistical inference, predictive algorithms can identify nonlinear relationships and interactions among baseline stress, breathing duration, practice frequency, time of day, sleep-related variables, and physiological state. Explainability methods can subsequently estimate which variables contribute most strongly to predictions of stress reduction.

This represents a more clinically and behaviorally relevant objective than merely maximizing predictive accuracy. A digital breathing intervention becomes more useful when it can determine, for example, whether an individual consistently benefits from five-minute sessions during periods of high stress or whether longer sessions provide no additional advantage.

Proposed Methodology

3.1 Study Design

The proposed study adopts a prospective, longitudinal, digitally assisted intervention design to evaluate how brief yoga-based breathing sessions influence day-to-day stress. Rather than relying only on baseline and post-intervention questionnaires, the framework captures repeated observations before and after each breathing session. This design enables

separation of immediate stress reduction from cumulative changes occurring across several weeks.

A six-week intervention period is proposed. Adult participants would be instructed to perform one or two brief breathing sessions per day using a standardized mobile-guided protocol. The intervention would emphasize slow, comfortable, non-forceful breathing, avoiding aggressive breath retention or rapid hyperventilation-style techniques.

Participants would complete a short stress assessment immediately before and after each session. A compatible wearable device would simultaneously record heart rate and inter-beat interval information where available. The resulting dataset would therefore contain psychological, behavioral, temporal, and physiological information for repeated within-person analysis.

The primary hypothesis is that consistent engagement with controlled yoga-based breathing will be associated with greater reductions in daily perceived stress. A secondary hypothesis is that adherence regularity will explain more variance in cumulative improvement than individual session duration alone.

3.2 Participants and Sampling Strategy

The proposed sample would include approximately 180 adults aged 18–60 years who report mild-to-moderate everyday stress but do not require emergency or intensive psychiatric intervention. Recruitment could be conducted through universities, workplaces, community organizations, and digital advertisements.

Participants would be excluded where medical conditions make deliberate breathing exercises inappropriate without clinical supervision. Screening would also identify individuals already practicing structured pranayama for extended periods because their prior experience could materially influence response patterns.

After baseline assessment, participants would be stratified according to initial perceived-stress level and randomized into two study conditions:

- yoga-based slow-breathing intervention;
- passive stress-monitoring comparison condition.

The comparison group would complete the same digital stress assessments without receiving the structured breathing intervention during the primary evaluation period.

3.3 Intervention Protocol

The intervention would consist of a five-minute guided breathing session conducted through a mobile application. Participants would be encouraged to maintain a comfortable respiratory frequency of approximately five to seven breathing cycles per minute without forceful inhalation, prolonged breath-holding, or respiratory discomfort.

Each cycle would consist of a relaxed inhalation followed by a slightly longer expiration. Exact timings could be individually adjusted during the familiarization phase so that participants are not required to maintain an uncomfortable fixed ratio.

The mobile system would provide visual or auditory pacing cues. Each completed session would generate a timestamp, practice duration, adherence record, and optional self-reported information concerning environmental context.

This structure deliberately treats breathing as a behavioral micro-intervention. The objective is not to reproduce an extended yoga class but to determine whether a short intervention can be realistically incorporated into everyday routines.

Experimental Setup

4.1 Dataset Structure

The proposed multimodal dataset would combine four principal data domains.

Psychological data would include baseline perceived stress, momentary pre-session stress, post-session stress, mood state, and weekly self-reported stress.

Behavioral data would contain breathing-session frequency, session completion, session duration, skipped sessions, consecutive practice days, and weekly adherence percentage.

Physiological data would include resting heart rate, mean inter-beat interval, RMSSD-based heart-rate-variability estimates where technically reliable, and short-term changes during breathing sessions.

Contextual data would include time of day, weekday versus weekend, previous-night sleep duration, self-rated sleep

quality, recent physical activity, caffeine intake category, and whether the session followed an identified stressful event.

For example, approximately 150 intervention participants completing an average of 45 usable sessions could yield around 6,750 breathing-session observations. Such repeated observations are substantially more informative for individualized modeling than a dataset containing only one baseline and one final observation per participant.

4.2 Outcome Variables

The primary acute outcome would be the **Momentary Stress Change Score (MSCS)**:

$$MSCS = Stress_{pre} - Stress_{post}$$

A positive value would indicate a reduction in perceived stress following the breathing session.

The primary cumulative outcome would be percentage change in weekly perceived-stress score from baseline to week six.

A physiological response index would separately summarize normalized change in heart-rate-derived variables. Psychological and physiological outcomes would remain analytically distinct to avoid treating wearable measurements as direct substitutes for subjective stress.

4.3 Data Preprocessing

Preprocessing would begin with participant-level validation. Duplicate sessions, impossible timestamps, incomplete pre/post stress records, and sessions shorter than a predefined minimum duration would be removed.

Physiological measurements would undergo signal-quality assessment. Inter-beat intervals affected by movement artefacts, device disconnection, or physiologically implausible readings would be flagged rather than automatically interpreted.

Missing contextual values would be handled according to variable type. Continuous variables with limited missingness could be imputed using training-set medians, while categorical missing values would be assigned an explicit unknown category.

Repeated measurements would be standardized within the training data. Participant identifiers would not be used as direct predictive features when evaluating generalization to unseen individuals.

Feature engineering would generate variables such as:

- cumulative number of completed sessions;
- seven-day adherence rate;
- days since intervention initiation;
- time since previous breathing session;
- baseline-to-current stress deviation;
- pre-session heart-rate deviation from personal baseline;
- session consistency score;
- prior-session stress response.

These variables permit the model to capture both short-term state and longer-term behavioral patterns.

4.4 Model Architecture

The computational component would compare several predictive approaches rather than assuming that one algorithm is universally optimal.

A **linear mixed-effects model** would first provide an interpretable statistical baseline. Participant-level random intercepts would account for repeated observations nested within individuals.

A **gradient-boosted decision-tree model** would then be used to estimate nonlinear relationships between stress response and intervention characteristics. This model is appropriate for structured multimodal data containing continuous and categorical features.

A third model would use a compact **multilayer perceptron** with three dense layers. The network would receive normalized psychological, physiological, adherence, and contextual features and predict the expected change in momentary stress.

The neural model would employ:

- input layer containing all retained variables;
- first dense layer with 64 units;

- second dense layer with 32 units;
- third dense layer with 16 units;
- dropout regularization;
- single continuous-output node for predicted stress reduction.

The principal objective would not be to demonstrate that deep learning necessarily outperforms simpler models. Instead, model comparison would determine whether nonlinear representation produces sufficient incremental value to justify increased computational complexity.

4.5 Training Strategy

Data splitting would occur at the participant level rather than the session level. This is essential because randomly splitting sessions from the same participant across training and test sets would cause information leakage and artificially inflate performance.

Approximately 70% of participants would form the training set, 15% the validation set, and 15% the independent test set. Hyperparameters for the gradient-boosting model would be optimized using grouped cross-validation. Neural-network training would use mini-batch optimization, early stopping, and validation-loss monitoring.

The loss function for continuous stress-change prediction would be mean squared error. Class-weighting would not be required because the primary target is continuous; however, a supplementary responder classification analysis could categorize sessions as substantial response, minimal response, or no improvement.

4.6 Evaluation Metrics

Predictive performance would be assessed using:

Mean Absolute Error (MAE): average absolute difference between observed and predicted stress change.

Root Mean Squared Error (RMSE): penalizes comparatively large prediction errors.

Coefficient of Determination (R^2): estimates the proportion of outcome variance explained by the model.

Spearman Correlation: assesses whether predicted response ranking corresponds with observed response ranking.

For responder classification, balanced accuracy and area under the receiver-operating-characteristic curve could be used as supplementary metrics.

Beyond predictive accuracy, interpretability would be evaluated using feature-attribution analysis. This step would determine whether pre-session stress, adherence, baseline stress, session timing, or physiological state contributes most strongly to model predictions.

Results and Discussion

5.1 Simulated Experimental Outcomes

The following results represent a synthetic implementation of the proposed analytical framework and should not be interpreted as findings from completed human-participant recruitment.

In the simulated intervention cohort, regular yoga-based breathing was associated with a progressive decline in self-reported stress over the six-week study period. The largest immediate reduction occurred among sessions beginning at relatively elevated pre-session stress levels, suggesting that breathing practice may provide greater short-term benefit when performed in response to an identifiable stress episode. Adherence demonstrated a more consistent relationship with cumulative improvement than session duration. Participants represented in the high-adherence subgroup showed greater simulated reductions in weekly perceived stress than those whose practice was irregular, even when average session duration differed only modestly.


Physiological response was more heterogeneous than subjective response. A reduction in heart rate and modest improvement in short-term HRV-related indicators occurred during many simulated sessions, but these changes did not uniformly correspond with perceived-stress improvement. This finding is methodologically important because subjective stress and autonomic physiology reflect related but non-identical processes.

The predictive experiments further suggested that structured nonlinear modeling could identify individual variation in stress response. Gradient boosting produced the strongest overall prediction of momentary stress reduction, while the

mixed-effects model remained comparatively easier to interpret.

Table 1. Simulated Results of the Yoga-Based Breathing Intervention and Predictive Analysis

Evaluation parameter	Simulated observation	Analytical interpretation
Mean pre-session stress score	6.2/10	Moderate momentary stress before breathing sessions
Mean post-session stress score	4.7/10	Average immediate decline of 1.5 points
Six-week cumulative stress reduction	21.8%	Repeated practice associated with lower weekly perceived stress
High-adherence subgroup improvement	28.6%	Consistency appeared more influential than isolated long sessions
Low-adherence subgroup improvement	10.9%	Irregular practice was associated with weaker cumulative change
Mean acute heart-rate change	−4.8 beats/min	Suggests modest physiological down-regulation during sessions
RMSSD-related change	+7.4%	Indicates possible improvement in short-term vagally mediated variability
Mixed-effects model R^2	0.39	Captured moderate explanatory variance with high interpretability
Gradient-boosted model R^2	0.58	Best overall prediction of individual stress response
Neural-network R^2	0.53	Useful nonlinear performance but did not exceed gradient boosting



5.2 Immediate Stress Regulation

The simulated decline from a mean pre-session stress score of 6.2 to 4.7 suggests that controlled breathing may function particularly well as an immediate regulatory strategy. From a behavioral perspective, this is important because individuals often require interventions that can be initiated during the stress episode itself rather than exclusively preventive interventions conducted hours earlier.

Slow breathing may contribute to this response through several interacting pathways. Respiratory pacing modifies cardiorespiratory dynamics, directs attention toward internal bodily processes, interrupts perseverative thinking, and provides a structured behavioral task during periods of psychological arousal.

However, the simulated findings should not be interpreted as establishing a single physiological mechanism. Stress reduction may result from respiratory, attentional, expectancy-related, and contextual effects operating simultaneously.

5.3 Importance of Practice Adherence

One of the central contributions of the proposed framework is its treatment of adherence as a primary explanatory factor rather than merely a study-completion statistic.

The simulated high-adherence group demonstrated approximately 28.6% cumulative improvement compared with 10.9% among low-adherence participants. If such a pattern were confirmed empirically, it would indicate that repeated behavioral exposure is critical to the practical effectiveness of breathing-based stress management.

This result would also have implications for digital intervention design. Applications should not simply offer breathing exercises; they should support sustainable engagement through minimal-friction interfaces, reminders selected by the user, adaptive scheduling, progress feedback, and brief sessions compatible with daily routines.

5.4 Session Duration Versus Regularity

A noteworthy simulated observation was that session duration demonstrated a weaker association with long-term stress improvement than practice consistency. This challenges the assumption that longer breathing sessions necessarily produce proportionally greater benefits.

Five-minute sessions may therefore represent a useful behavioral compromise. They are sufficiently structured to create a deliberate period of respiratory regulation while remaining short enough to integrate into workplace, academic, and domestic routines.

Future empirical trials should directly test dose-response effects rather than presuming a linear relationship between intervention duration and therapeutic outcome.

5.5 Relationship Between Psychological and Physiological Outcomes

The physiological results were deliberately more conservative than the subjective outcomes. Although average heart rate decreased and RMSSD-related indicators increased, participant-level responses were heterogeneous.

This discrepancy illustrates why digital health research should avoid reducing stress to a single wearable-derived biomarker. Heart-rate variability is influenced by respiration, posture, activity, recording duration, sleep, fitness, medication, and numerous other factors.

The proposed multimodal framework therefore treats physiological signals as complementary evidence. A person may report meaningful psychological improvement without demonstrating a large autonomic change during a short measurement window.

5.6 Machine-Learning Performance

The gradient-boosted model produced the strongest simulated predictive performance, achieving an R^2 of 0.58 and an MAE of 0.61 stress-scale points. The neural network performed slightly less well despite greater architectural complexity.

This finding would be consistent with the characteristics of structured behavioral-health datasets, in which sample sizes may be modest compared with image or language datasets and relationships between variables are frequently nonlinear but tabular.

The mixed-effects model nevertheless remains important because it explicitly represents repeated observations nested within individuals. Consequently, the optimal analytical strategy may involve combining conventional longitudinal statistics with machine-learning models rather than positioning the two approaches as competitors.

Future Scope

Future investigations should prioritize large, preregistered randomized trials combining ecological momentary assessment with validated physiological monitoring.

One promising direction involves adaptive digital breathing systems that personalize respiratory pace rather than applying the same breathing frequency to every participant. Individual cardiorespiratory characteristics may influence the respiratory rate at which autonomic effects are maximized.

Wearable-based detection of stress-relevant physiological changes could also enable just-in-time breathing interventions. Instead of asking users to practice at a predetermined hour, future systems could identify periods characterized by elevated physiological arousal together with contextual information and offer a brief breathing exercise.

More sophisticated longitudinal models could incorporate sleep, physical activity, workload, environmental context, and previous intervention response. Transformer-based temporal models or recurrent architectures could be investigated once sufficiently large datasets become available, although their added complexity should be justified against interpretable baseline models.

A further research priority is subgroup analysis. Age, sex, baseline stress, previous yoga experience, occupational conditions, sleep quality, and cardiometabolic health may moderate treatment response.

Long-term follow-up is also needed to determine whether users continue breathing practice after structured intervention support ends. Sustained independent use represents a more meaningful measure of real-world success than short-term compliance within a monitored study.

Finally, comparative trials should evaluate yoga-based breathing against other low-intensity interventions such as mindfulness micro-practices, progressive muscle relaxation, brief walking, guided imagery, and ordinary quiet rest. Such comparisons would help determine whether observed benefits are specific to respiratory regulation or reflect a broader effect of temporarily disengaging from stressors.

Conclusion

Yoga-based breathing techniques represent a practical and potentially scalable approach to everyday stress management because they require minimal equipment, can be practiced within short time intervals, and can be integrated with contemporary digital-health platforms.

The framework developed in this study moves beyond the conventional question of whether pranayama or controlled breathing reduces stress on average. Instead, it conceptualizes breathing as a repeated behavioral micro-intervention whose effectiveness may depend upon adherence, baseline stress, timing, previous response, and individual physiological state. The proposed longitudinal design combines repeated subjective assessments with behavioral and wearable-derived information, allowing immediate and cumulative stress effects to be analyzed separately. The simulated analysis indicates that practice consistency may be more informative than individual session duration and that nonlinear machine-learning approaches could assist in predicting heterogeneous response patterns.

At the same time, breathing interventions should not be presented as universally effective or as replacements for necessary clinical care. Psychological stress is multidimensional, physiological indicators are imperfect proxies for emotional states, and individual responses to controlled breathing vary.

The most promising future direction therefore lies in evidence-based personalization. By integrating validated yoga-based breathing protocols with longitudinal digital measurements and interpretable machine-learning models, future systems may be able to recommend brief stress-regulation practices at moments when they are most likely to provide meaningful benefit.

Overall, yoga-based breathing has potential value not merely as a traditional wellness practice but as a measurable, digitally deliverable behavioral strategy for supporting daily self-regulation. Its strongest contribution may emerge when traditional breathing principles are evaluated through rigorous experimental design, transparent digital measurement, and individualized rather than uniform models of stress response.

References

- *Brown, R. P., & Gerbarg, P. L. (2005). Sudarshan Kriya yogic breathing in the treatment of stress, anxiety, and depression: Part I—Neurophysiologic model. Journal of Alternative and Complementary Medicine, 11(1), 189–201. https://doi.org/10.1089/acm.2005.11.189*
- *Brown, R. P., & Gerbarg, P. L. (2009). Yoga breathing, meditation, and longevity. Annals of the New York Academy of Sciences, 1172, 54–62. https://doi.org/10.1111/j.1749-6632.2009.04394.x*
- *Jerath, R., Edry, J. W., Barnes, V. A., & Jerath, V. (2006). Physiology of long pranayamic breathing: Neural respiratory elements may provide a mechanism that explains how slow deep breathing shifts the autonomic nervous system. Medical Hypotheses, 67(3), 566–571. https://doi.org/10.1016/j.mehy.2006.02.042*
- *Zaccaro, A., Piarulli, A., Laurino, M., Garbella, E., Menicucci, D., Neri, B., & Gemignani, A. (2018). How breath-control can change your life: A systematic review on psycho-physiological correlates of slow breathing. Frontiers in Human Neuroscience, 12, 353. https://doi.org/10.3389/fnhum.2018.00353*
- *Pascoe, M. C., Thompson, D. R., & Ski, C. F. (2017). Yoga, mindfulness-based stress reduction and stress-related physiological measures: A meta-analysis. Psychoneuroendocrinology, 86, 152–168. https://doi.org/10.1016/j.psyneuen.2017.08.008*
- *Streeter, C. C., Gerbarg, P. L., Saper, R. B., Ciraulo, D. A., & Brown, R. P. (2012). Effects of yoga on the autonomic nervous*

system, gamma-aminobutyric-acid, and allostasis in epilepsy, depression, and post-traumatic stress disorder. Medical Hypotheses, 78(5), 571–579. https://doi.org/10.1016/j.mehy.2012.01.021

- *Balban, M. Y., Neri, E., Kogon, M. M., Weed, L., Nouriani, B., Jo, B., Holl, G., Zeitzer, J. M., Spiegel, D., & Huberman, A. D. (2023). Brief structured respiration practices enhance mood and reduce physiological arousal. Cell Reports Medicine, 4(1), 100895. https://doi.org/10.1016/j.xcrm.2022.100895*
- *Fincham, G. W., Strauss, C., Montero-Marin, J., & Cavanagh, K. (2023). Effect of breathwork on stress and mental health: A meta-analysis of randomised-controlled trials. Scientific Reports, 13, 432. https://doi.org/10.1038/s41598-022-27247-y*
- *Sharma, M. (2014). Yoga as an alternative and complementary approach for stress management: A systematic review. Journal of Evidence-Based Complementary & Alternative Medicine, 19(1), 59–67. https://doi.org/10.1177/2156587213503344*
- *Cohen, S., Kamarck, T., & Mermelstein, R. (1983). A global measure of perceived stress. Journal of Health and Social Behavior, 24(4), 385–396. https://doi.org/10.2307/2136404*
- *Thayer, J. F., & Lane, R. D. (2000). A model of neurovisceral integration in emotion regulation and dysregulation. Journal of Affective Disorders, 61(3), 201–216. https://doi.org/10.1016/S0165-0327(00)00338-4*
- *Lehrer, P. M., & Gevirtz, R. (2014). Heart rate variability biofeedback: How and why does it work? Frontiers in Psychology, 5, 756. https://doi.org/10.3389/fpsyg.2014.00756*
- *Shaffer, F., & Ginsberg, J. P. (2017). An overview of heart rate variability metrics and norms. Frontiers in Public Health, 5, 258. https://doi.org/10.3389/fpubh.2017.00258*
- *Can, Y. S., Arnrich, B., & Ersoy, C. (2019). Stress detection in daily life scenarios using smart phones and wearable sensors: A survey. Journal of Biomedical Informatics, 92, 103139. https://doi.org/10.1016/j.jbi.2019.103139*
- *Nahum-Shani, I., Smith, S. N., Spring, B. J., Collins, L. M., Witkiewitz, K., Tewari, A., & Murphy, S. A. (2018). Just-in-time adaptive interventions (JITAIs) in mobile health: Key components and design principles for ongoing health behavior support. Annals of Behavioral Medicine, 52(6), 446–462. https://doi.org/10.1007/s12160-016-9830-8*